






ABOVE: KILLDEER (*Charadrius vociferus*). Summer: southern Canada and northern U.S. Year-round: southern U.S. BELOW: MALLARD (*Anas platyrhynchos*). Female in foreground, male in background. Summer: most of Canada. Winter: most of U.S.





SLATE-COLORED JUNCO (*Junco hyemalis*). Male in foreground, female in background. Summer: most of Canada. Winter: most of U.S.



RUBY-THROATED HUMMINGBIRD (*Archilochus colubris*). Male left, female right. Found in summer in eastern half of U.S.






GOLDFINCH (Common Goldfinch; American Goldfinch; *Spinus tristis*).
Female in center, others male. Found throughout U.S. with seasonal range
variation.





CATBIRD (*Dumetella carolinensis*). Summer: most of U.S. except West Coast. Winter: extreme southeastern U.S.



DOWNY WOODPECKER (*Dendrocopus pubescens*). Female left, male right. Year-round in most of U.S. and wooded parts of Canada.




CARDINAL (*Richmondena cardinalis*). Female left, male right. Year-round in most parts of the U.S.





BLUE JAY (*Cyanocitta cristata*). Found year-round in eastern two-thirds of U.S. and southeastern Canada.



ABOVE: ROBIN (*Turdus migratorius*). Summer: most of Canada. Year-round: most of U.S. BELOW: WOOD THRUSH (*Hyclocichla ustulata*). Found in summer in eastern half of U.S.





SPARROW HAWK (*Falco sparverius*). Summer: most of Canada, northern U.S. Year-round: southern U.S. and West Coast.





RED-HEADED WOODPECKER (*Melanerpes erythrocephalus*). Summer: north central U.S. Year-round: most of eastern U.S.



BALTIMORE ORIOLE (*Icterus galbula*). Male above, female below. Found in summer in southeastern Canada and eastern two-thirds of the U.S.










visit the world's
SUPER
COLORING





Printed in the USA
SUPER
COLORING



Join the world's
SUPER
COLORING



print the world
SUPER
COLORING

